

(19) World Intellectual Property Organization
International Bureau(43) International Publication Date
30 May 2002 (30.05.2002)

PCT

(10) International Publication Number
WO 02/42290 A1(51) International Patent Classification⁷: C07D 401/04,
A61K 31/445

(21) International Application Number: PCT/HU01/00111

(22) International Filing Date:
14 November 2001 (14.11.2001)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
P 0004701 23 November 2000 (23.11.2000) HU(71) Applicant (for all designated States except US):
RICHTER GEDEON VEGYÉSZETI GYÁR RT.
[HU/HU]; Gyömroi út 19-21, H-1103 Budapest (HU).

(72) Inventors; and

(75) Inventors/Applicants (for US only): FISCHER, János
[HU/HU]; Úri u. 33, H-1014 Budapest (HU). FODOR,
Tamás [HU/HU]; Andrásy ut 3, H-1061 Budapest (HU).
TRISCHLER, Ferenc [HU/HU]; Úttörő u. 16, H-1171
Budapest (HU). LÉVAI, Sándor [HU/HU]; Ipar u. 20,
H-2051 Biantorbágy (HU). PETÉNYI, Endréné [HU/HU];
Róbert Károly krt. 16/C., H-1138 Budapest (HU).(74) Common Representative: RICHTER GEDEON VEG-
YÉSZETI GYÁR RT.; Gyömroi út 19-21, H-1103 Bu-
dapest (HU).(81) Designated States (national): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM,
HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK,
LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX,
MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL,
TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.(84) Designated States (regional): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian
patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European
patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE,
IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF,
CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD,
TG).

Declaration under Rule 4.17:


— as to applicant's entitlement to apply for and be granted a patent (Rule 4.17(ii)) for the following designations AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG)

Published:


— with international search report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: NEW DESLORATADINE SALTS, PROCESS FOR THEIR SYNTHESIS AND PHARMACEUTICAL COMPOSITIONS THEREOF



H-X (I)



2 H-X (II)

(57) Abstract: The object of the present invention are new desloratadine salts of formula I wherein the meaning of X is an acid residue and the meaning of n is 1 or 2, and formula II wherein the meaning of X is a pK < 3.5 acid residue. The invention is related to a process for their synthesis, as well as new anti-allergic pharmaceutical compositions containing these salts.

New desloratadine salts, process for their synthesis and pharmaceutical compositions thereof

The invention relates to new desloratadine salts, process for their synthesis, as well
5 as new anti-allergic pharmaceutical compositions containing these salts.

It is known, that desloratadine (it's chemical name: 8-chloro-6,11-dihydro-11-(4-piperidilydene)-5H-benzo[5,6]cyclohepta[1,2-b]pyridine) is an active metabolite of a
10 successful anti-allergic drug substance, loratadine. According to the literature desloratadine is 2.5-4 times more active orally than loratidine and antihistaminic activity lasts for 24 h (Arzneim. Forsch./Drug Res. 50 (I), Nr. 4 (345-352) 2000).


It is known from the Hungarian patent Number 194 864, that desloratadine base can
be obtained from loratadine (chemical name: 8-chloro-6,11-dihydro-11-(1-ethoxycarbonyl-4-piperidilydene)-5H-benzo[5,6]cyclohepta[1,2-b]pyridine) by two
15 methods. These are as follows:

a) the 8-chloro-6,11-dihydro-11-(1-ethoxycarbonyl-4-piperidilydene)-5H-benzo
[5,6]cyclohepta[1,2-b]pyridine (loratadine) is decarbethoxylated by boiling with
20 aqueous ethanolic sodium hydroxide solution for 24 h, then isolating the desloratadine acetate after neutralizing the solution with acetic acid. This crude product has to be further purified; the desloratadine acetate - according to the paper - is obtained in 70 % yield after recrystallization from benzene-hexane mixture. The desloratadine base is prepared by basic treatment of desloratadine acetate and this
25 is purified by recrystallization from hexane.

b) the 8-chloro-6,11-dihydro-11-(1-methyl-4-piperidilydene)-5H-benzo[5,6]
cyclohepta[1,2-b]pyridine is demethylated in two steps: first the 1-cyano-derivative is
synthesized with cyanogen bromide and this is hydrolyzed with concentrated
30 hydrochloric acid solution in acetic acid for 20 h, then after evaporating the solvents the residue is neutralized with ammonium hydroxide solution to obtain the desloratadine, the melting point of which is 149-151 °C.


It is mentioned in the above Hungarian patent, that salts can be formed from desloratadine with pharmaceutically acceptable acids: hydrochloric acid, methanesulfonic acid, sulfuric acid, acetic acid, maleinic acid, fumaric acid, phosphoric acid, but the formula, the physical- and physicochemical data and the
5 method of their synthesis – except the above acetate salt – are not given.

- The above mentioned processes for the synthesis of desloratadine have several
disadvantages. During the realization of process a) substantial decomposition takes
place, therefore, there are several impurities in the final product. The desloratadine
10 base of required purity can be obtained by recrystallization, but this process can be
carried out only with substantial loss of material. During the formulation of the active
ingredient considerable disadvantage is from the point of technology, that the
desloratadine base is insoluble in water.
- 15 Process b) is disadvantageous in itself, because of the use of poisonous cyanogen
bromide reagent and the poisonous methyl bromide formed in the two-step reaction.
On the other hand, the desloratadine base obtained by the latter method has the
same disadvantages as the one obtained by method a).
- 20 In our experiments surprisingly we found that desloratadine acid addition salts of
formula I



3

wherein the meaning of X is halogen atom, preferably chlorine or bromine, or acid residue, the meaning of n is 1 or 2, can be obtained by treatment/heating of loratadine base of formula III with certain acids.



(III)


5

The so obtained acid addition salts are new and among them the desloratadine hemisulfate is particularly advantageous, because it can be obtained in one step, in high purity and stability. The other properties of the new acid addition salts are also favorable, for example their good solubility is advantageous from the point of drug formulation.

10

According to the above mentioned facts the invention relates to acid addition salts of formula I - wherein the meaning of X is an acid residue and the meaning of n is 1 or 2 - as well as the acid addition salts of formula II

15



(II)

- wherein the meaning of X is an acid residue of $pK < 3.5$ acid.

The invention also relates to the synthesis of acid addition salts of formula II, by reacting the loratadine of formula III (chemical name: 8-chloro-6,11-dihydro-11-(1-ethoxycarbonyl-4-piperidylidene)-5H-benzo[5,6]cyclohepta[1,2-b]pyridine) with concentrated mineral acid.

5

Further object of the invention is the method for the synthesis of acid addition salts of formula I – wherein the meaning of X is an acid residue and the meaning of n is 1 or 2 – by treating an acid addition salt of formula II – wherein the meaning of X is an acid residue of $pK < 3.5$ acid – or an aqueous solution of it with a solution of a base to adjust the pH to 6.5-7, then isolating the product.

10

Finally the invention relates to anti-allergic pharmaceutical composition containing 0.1-99.9 % of active ingredient of formula I or II and 0.1-99.9 % of pharmaceutically acceptable carriers and additives.

15

Detailed description of the process:

20

In the process according to our invention the loratadine is heated with concentrated mineral acids, this way the urethane is hydrolyzed in a few hours and the salt of desloratadine formed with two mole acid (see formula II, wherein the meaning of X is as given above) can be isolated in good yield.

25

According to a preferred realization of the invention the loratadine is heated with 60-80 wt. % sulfuric acid solution at 110-120 °C, this way the hydrolysis of the urethane takes 3-6 h. The desloratadine disulfate can be isolated from the reaction mixture in good yield (80-95 %).

30

According to an other preferred realization of the invention the loratadine is heated with concentrated hydrochloric acid at 115 °C, this way the hydrolysis of the urethane takes 6 h and the desloratadine dihydrogen chloride salt can be isolated from the reaction mixture in high yield (90-95 %).

According to a further realization of the invention the loratadine is heated with 48 % hydrogen bromide solution at 110 °C. This way the urethane is hydrolyzed in 6 h and the desloratadine dihydrogen bromide salt can be isolated in high yield (> 95 %).

- 5 The desloratadine double salts can be isolated not only in good yield, but in high purity as well.

According to our invention the desloratadine double salts can be transformed into simple salts with strong base.

10

Especially preferred the formation of desloratadine hemisulfate from desloratadine disulfate with addition of strong base, for example 25 % tetramethylammonium hydroxide solution, to adjust the pH to 6.8 and isolating the desloratadine hemisulfate.

15

The new desloratadine hemisulfate of our invention can be the active ingredient of a new, non-sedative H1-antagonist pharmaceutical composition.

20

The starting material of the compounds of the invention is loratadine (chemical name: 8-chloro-6,11-dihydro-11-(1-ethoxycarbonyl-4-piperidylidene)-5H-benzo[5,6]cyclohepta[1,2-b]pyridine). The synthesis of loratadine is described in detail in the US patent Number 4 282 233 (the equivalent of which is the Hungarian patent Number 186 774).

- 25 The invention is illustrated by the following not limiting Examples:

Example 1

Desloratadine disulfate

30

A mixture of 19.5 g (50 mmol) of loratadine and 40 g of 72 wt. % sulfuric acid is stirred at 115 °C for 6 h. The reaction mixture is cooled to room temperature, 100 ml

of methanol is added, then the mixture is cooled to 0 °C and stirred at this temperature for 3 h. The precipitated crystalline product is filtered off, washed with ice-cold methanol. After drying 20.95 g (84 %) of the title compound is obtained. Melting point: 244-246 °C.

- 5 According to HPLC measurements the purity of the product is > 99.5 %.

Determination by titrimetry:

- 10 The desloratadine disulfate is dissolved in 80 % acetone and it is titrated with 0.1 N sodium hydroxide solution by potentiometry. The titration curve has two inflection points; the two bisulfate anion and the proton on the nitrogen of the pyridine are titrated till the first inflection point and the proton on the nitrogen of the piperidine is titrated between the two inflections. The ratio of the two area is 3/1.

15

Example 2

Desloratadine dihydrogen chloride

- 20 A mixture of 5.0 g (13 mmol) of loratadine (in solid form) and 50 ml of concentrated hydrochloric acid is stirred at 115 °C for 6 h. The excess of hydrochloric acid is evaporated and the residue is crystallized with 30 ml of acetone. The crystalline suspension is stirred at 0 °C for 5 h, filtered and washed with acetone to yield 4.7 g (94%) of the title compound. Melting point: 210-220 °C.

25

Example 3

Desloratadine dihydrogen bromide

30

A mixture of 3.83 g (10 mmol) of loratadine and 30 ml of 48 % hydrogen bromide is stirred at 115 °C for 6 h. The excess of hydrogen bromide is evaporated and the

residue is dissolved in 20 ml of hot ethanol. The title compound is precipitated in crystalline form after cooling. The crystalline suspension is stirred at 0 °C for 3 h, filtered and washed with ice-cold ethanol to yield 4.7 g (99 %) of the title compound. Melting point: 247-250 °C.

5

Example 4

Desloratadine hemisulfate

10

3.04 g (6 mmol) of desloratadine disulfate (obtained according to Example 1) is dissolved in a mixture of 5 ml of water and 2 ml of ethanol, then cooled to 0 °C and the pH is adjusted to 6.8 with addition of 25 % tetramethylammonium hydroxide solution. The solvent is evaporated and the residue is stirred with 50 ml of ethanol at 0 °C for 5 h, filtered and washed with ice-cold ethanol to yield 1.64 g (76 %) of the title compound. Melting point: 279-280 °C.

15

Determination by titrimetry:

The desloratadine hemisulfate is dissolved in 80 % acetone and it is titrated with 0.1 N sodium hydroxide solution by potentiometry. Only one inflection point is observed, which is equivalent with the proton on the nitrogen of the piperidine.

20

Example 5

25

General procedure for the preparation of salts of formula I

5.07 g (10 mmol) of desloratadine disulfate is suspended in 50 ml of dichloromethane and 10 ml of 4N sodium hydroxide solution is added. After vigorous stirring the solutions clear up. The organic layer is separated, washed with 10 ml of saturated sodium chloride solution and dried over anhydrous magnesium sulfate.

30

10 mmol of acid of formula HX is added to the dichloromethane solution. The product is precipitated from the solution in crystalline form after cooling.

The following salts of formula I were prepared:

n	X	Melting point (°C)	pH of 1 % solution	H ₂ O	Yield (%)
1	C ₆ H ₅ -SO ₃	212-214	5.6	0	91
1	$\begin{array}{c} \text{CH}_2\text{-COOH} \\ \\ \text{HO-C-COOH} \\ \\ \text{CH}_2\text{-COO} \end{array}$ <i>maleic acid</i>	63-114	4.5	2	95
1	$\begin{array}{c} \text{COO} \\ \\ \text{CH-OH} \\ \\ \text{CH-OH} \\ \\ \text{COOH} \end{array}$	183	4.2	2	99
1	CH ₃ -SO ₃ <i>sulfonic acid</i>	242-247	5.2	0	95
1	HSO ₄ <i>sulfuric acid</i>	237-247	3.0	0	88
1	Cl <i>hydrochloric acid</i>	271-273	4.8	0	77
1	$\begin{array}{c} \text{CH-COOH} \\ \\ \text{CH-COO} \end{array}$	169-171	5.0	0	94

5

maleic acid

Example 6

Preparation of a pharmaceutical composition

10 For 100 mg tablets the following ingredients are required (for one tablet):

desloratadine hemisulfate 5.0 mg

(prepared according to Example 4)

lactose 47.0 mg

9

corn-starch 47.0 mg

magnesium stearate 1.0 mg

The mixture of the powders is pressed into tablets directly after homogenization.

5

Example 7

Preparation of a pharmaceutical composition

For 100 mg tablets the following ingredients are required (for one tablet):

10 desloratadine hemisulfate 5.0 mg
(prepared according to Example 4)
lactose 25.0 mg
corn-starch 69.0 mg
magnesium stearate 1.0 mg

15 The mixture of the powders is pressed into tablets directly after homogenization.

Example 8

20 Preparation of a pharmaceutical composition


For 100 mg tablets the following ingredients are required (for one tablet):

desloratadine hemisulfate 5.0 mg
(prepared according to Example 4)
lactose 69.0 mg
25 corn-starch 25.0 mg
magnesium stearate 1.0 mg

The mixture of the powders is pressed into tablets directly after homogenization.


What we claim is:

1. The acid addition salts of formula I



- wherein the meaning of X is an acid residue and the meaning of n is 1 or 2.


2. The acid addition salts of formula II



- wherein the meaning of X is a $pK < 3.5$ acid residue.

3. Process for the synthesis of acid addition salts of formula II, characterized by reacting the loratadine of formula III

11



(III)

(chemical name: 8-chloro-6,11-dihydro-11-(1-ethoxycarbonyl-4-piperidylidene)-5H-benzo[5,6]cyclohepta[1,2-b]pyridine) with concentrated mineral acid.

4. Process for the synthesis of acid addition salts of formula I – wherein the meaning of X is an acid residue and the meaning of n is 1 or 2 – characterized by treating an acid addition salt of formula II – wherein the meaning of X is a $pK_a < 3.5$ acid residue - or the aqueous solution thereof with a solution of a base to adjust the pH to 6.5-7 and isolating the product.
5. Anti-allergic pharmaceutical composition, characterized by containing 0.1-99.9 % of active ingredient of formula I or II and 0.1-99.9 % of pharmaceutically acceptable carriers and additives.

INTERNATIONAL SEARCH REPORT

Int. Application No

PCT/HU 01/00111

A. CLASSIFICATION OF SUBJECT MATTER
IPC 7 C07D401/04 A61K31/445

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (Classification system followed by classification symbols)

IPC 7 C07D A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	EP 0 152 897 A (SCHERING CORP) 28 August 1985 (1985-08-28) cited in the application examples I, II, III	1-5
A	US 4 282 233 A (VILANI FRANK J) 4 August 1981 (1981-08-04) cited in the application example 1B	1-5

☐ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents :

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

T later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

X document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

Y document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

G document member of the same patent family

Date of the actual completion of the international search

31 January 2002

Date of mailing of the international search report

07/03/2002

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2200 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Bakboord, J

INTERNATIONAL SEARCH REPORT

Int Application No

PCT/HU 01/00111

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
EP 0152897	A	28-08-1985	AT 47140 T	15-10-1989
			AU 570306 B2	10-03-1988
			AU 3993885 A	10-09-1985
			DE 3573600 D1	16-11-1989
			DK 469485 A ,B,	14-10-1985
			EP 0152897 A1	28-08-1985
			FI 853675 A ,B,	25-09-1985
			HU 38332 A2	28-05-1986
			JP 5072910 B	13-10-1993
			JP 61501205 T	19-06-1986
			LU 90738 A9	09-05-2001
			WO 8503707 A1	29-08-1985
			US 4659716 A	21-04-1987
<hr/>				
US 4282233	A	04-08-1981	AT 9695 T	15-10-1984
			AU 543054 B2	28-03-1985
			AU 7186281 A	24-12-1981
			CA 1160230 A1	10-01-1984
			CY 1405 A	22-04-1988
			DE 3166441 D1	08-11-1984
			DK 263481 A	20-12-1981
			EP 0042544 A2	30-12-1981
			ES 503085 D0	01-11-1982
			ES 8300779 A1	01-02-1983
			FI 811878 A ,B,	20-12-1981
			HK 94387 A	18-12-1987
			HU 186774 B	30-09-1985
			IE 51303 B1	26-11-1986
			IL 63122 A	30-06-1985
			JP 1506964 C	13-07-1989
			JP 57035586 A	26-02-1982
			JP 63055513 B	02-11-1988
			KE 3758 A	02-10-1987
			KR 8500744 B1	24-05-1985
			LU 88359 A9	04-05-1994
			MY 76187 A	31-12-1987
			NZ 197435 A	30-03-1984
			PH 19252 A	17-02-1986
			PT 73200 A ,B	01-07-1981
			SG 70587 G	19-02-1988
			US 4355036 A	19-10-1982
			US 4560688 A	24-12-1985
			US 4831042 A	16-05-1989
			ZA 8104062 A	28-07-1982